

Looking Down the Light Cone: Can Deep Redshift Surveys Alone Measure the Power Spectrum?

Andrew A. de Laix and Glenn D. Starkman

Physics Department, Case Western Reserve University, Cleveland, OH 44106-7079

Received _____; accepted _____

ABSTRACT

We analyze the window functions for the spherical harmonic mode estimators of all-sky, volume limited surveys considering evolutionary effects along the past light-cone which include the deviation of the distance scale from a linear relationship with redshift, linear peculiar velocity corrections, and linear evolution of the density perturbations. The spherical harmonic basis functions are considered because they correspond most closely to the symmetries of typical survey geometries and of the light-cone effects we consider. Our results show substantial broadening of the windows over that expected by ignoring light-cone effects, indicating the difficulty of measuring the power spectrum independently from cosmology. We suggest that because of light-cone effects, deep redshift surveys should either be analyzed in conjunction with CMBR data which determines the cosmological parameters, or by using a Bayesian likelihood scheme in which varying cosmological parameters and a simple parameterization of the primordial power spectrum are assumed as the priors, so that observed data can be mapped from redshift to real space. The derived power spectrum can then be compared to underlying models of fluctuation generation and growth in structure formation to evaluate both these models and the cosmological priors.

The ability of a number of current and proposed galaxy surveys, *e.g.* The Sloan Digital Sky Survey (hereafter SDSS, Gunn & Weinberg (1995), Strauss (1997)) and the two-degree-field survey (Taylor & Gray (1994)), to probe deep redshifts offers exciting opportunities to observe structure on the largest scales, yielding results which will be invaluable for cosmology. As is often the case, however, the data collected is not directly compatible with that which theorists calculate. To see this, consider a typical theoretical calculation of large scale structure. Structure is presumed to have evolved from the gravitational collapse of small initial fluctuations, and on scales larger than approximately 10 Mpc, this collapse is well described by linear perturbation theory. Usually, theorists express the outcomes of their structure-formation models in terms of the power spectrum, the Fourier transforms of the two-point correlation function given in conformal spatial coordinates, evaluated on a constant time slice. That is, they measure the power spectrum of the structure for a snapshot of the universe taken at a fixed time. A galaxy survey, however, detects structure on a light-cone time slice, and measures redshifts, not conformal distances. For lengths much shorter than the Hubble length, the distinction is unimportant since to first order distance is proportional to redshift, and structure does not evolve much over the light-crossing time of the survey. But, the newest surveys, such as the SDSS, will penetrate to as much as 50% of the Hubble distance. Either assuming that proper distance is linearly proportional to redshift over these distance scales, or ignoring the evolution of structure over the corresponding time scales, leads to significant errors in the estimation of the power spectrum. In this letter, we examine these effects and determine their significance for estimating the power spectrum from a deep galaxy survey.

All surveys are limited in the volume of space which they can cover, and so are less than ideal observations. Furthermore, shot noise arises from the finite sampling inherent in any galaxy survey. Conversely, theoretical predictions for power spectra are determined for the whole universe continuously sampled and are therefore expressed in terms of a continuous power spectrum (unless the universe is spatially compact, but no current evidence suggests that it is). In sampling a finite volume, one probes instead a discrete set of eigenmodes and the amplitudes of the observed modes will be a convolution of the continuum modes with a window function. Ideally, this window function is narrow so that the observed mode amplitudes correspond closely to their continuum counterparts, and this can be achieved through the proper weighting of the observational data.

For Fourier decomposition, Tegmark (1995) has developed a method of analyzing redshift surveys

guaranteed for a specified survey geometry to minimize the width of the window function to the shot noise limit. Fourier decomposition, however, is not well suited to the analysis of most redshift surveys because survey geometries are usually defined by slices of a sphere, and because redshift effects depend only on the distance from the observer. A better basis is one which takes advantage of this symmetry, *i.e.* the spherical harmonic basis (Heavens and Tabor (1995)). For standard homogeneous, isotropic models, the power spectrum of the spherical harmonic decomposition is equivalent to that of the Fourier decomposition — each depends only on the magnitude of the mode considered, so theoretical calculations are equally useful for each basis. Our analysis therefore addresses the determination or estimation of the power spectrum in the spherical harmonic basis.

Given a survey geometry, we would like to find a weighting function ψ which will produce the “optimal” estimate for the amplitude of a given ℓ_0, m_0 , and k_0 harmonic of the continuum fluctuation spectrum. By optimal we mean that ψ should simultaneously minimize the width of the window function in ℓ, m , and k space and minimize the shot noise signal due to finite sampling—under the constraint that ψ is non-trivial. Following Tegmark (1995), we define the penalty function

$$|\ell - \ell_0|^2 + |m - m_0|^2 + \frac{|k^2 - k_0^2|^2}{\gamma^2}, \quad (1)$$

which forces our window toward the ℓ_0, m_0 , and k_0 we are trying to observe. The shot noise is proportional to $n(z)^{-1}$, where $n(z)$ is the average number of galaxies observed in a unit volume at redshift z . To find the optimal ψ we solve the eigen value problem

$$\left(|\hat{\ell} - \ell_0|^2 + |\hat{m} - m_0|^2 + \frac{|\hat{k}^2 - k_0^2|^2}{\gamma^2} + \frac{1}{n(z)} \right) \psi = E\psi, \quad (2)$$

where the ground state represents the optimal weighting function as it minimizes E . We normalize it, for convenience, by requiring

$$\int d^3z \psi^2(z) = 1. \quad (3)$$

Note that ψ can always be chosen to be a real valued function (which we shall assume it is), simplifying the numerics. Also note that for the radial portion of the penalty function we choose $|\hat{k}^2 - k_0^2|^2$ instead of $|\hat{k} - k_0|^2$, which one might naively select. We do so because there is no simple way to express $\hat{k}$ in coordinate space; the operator $\hat{k}^2$ is simply negative the Laplacian in physical space. The operators $\hat{\ell}$ and $\hat{m}$ return the magnitude of the total angular momentum and the z component respectively. Since the angular portion is separable, we can re-express ψ as a product $\psi_{\mu\nu k}(z)Y_{\mu\nu}(\Omega_z)$ where μ and ν are the eigenvalues of the generalized spherical harmonics which satisfy the boundary conditions given by the survey geometry.

In the limit of an all sky survey, $\mu = \ell_0$ and $\nu = m_0$ which are integers, although this not generally so. The parameter γ controls how we wish to weight shot noise to the window width. By increasing γ , we can reduce shot noise by accepting a larger variation in k , or by reducing γ we can get a narrower k variation, but at the expense of increased shot noise. For the special case of a volume limited survey in which $n(z)$ is a constant, the shot noise can be absorbed into the right hand side of the equation and the choice of γ is irrelevant; there is only one optimal ψ . As we have already pointed out, the angular part of the solution is given by the generalized spherical harmonics. The solution to the radial portion is then just a spherical Bessel function $j_\mu(k_\mu z)$, where $j_\mu(k_\mu z_{max}) = 0$ at the outer boundary of the volume z_{max} , and k_μ is the zero closest to k , the wavenumber of the mode whose amplitude we wish to estimate. It is this simplified problem which we will consider in this letter since the numerics are easier, and we expect that any light-cone effects which we observe in a more general survey of similar depth will be at least if not more severe.

Given the optimal weighting function, our estimator for the fluctuation amplitude of the k, ℓ, m mode is

$$\hat{a}_{lm}(k) = \int d^3z \frac{n(\vec{z}) - \bar{n}(\vec{z})}{\bar{n}(\vec{z})} \psi(\vec{z}), \quad (4)$$

where $\bar{n}(\vec{z}) = \langle n(\vec{z}) \rangle$ is the ensemble average of the number density of galaxies at redshift z . A more interesting quantity is the square of this estimator, which averaged over the ensemble of realizations should be related to the continuum power spectrum. Using the analysis of Peebles (1980), one can straightforwardly show that in the ensemble average

$$\langle \hat{a}_{lm}(k) \hat{a}_{lm}(k) \rangle = \int d^3z \frac{\psi^2(\vec{z})}{\bar{n}(\vec{z})} + \int d^3z d^3z' \xi(\vec{z}, \vec{z}') \psi(\vec{z}) \psi(\vec{z}'), \quad (5)$$

where ξ is the correlation function in redshift space. From the above, we can infer an estimator for the power in a particular mode

$$\hat{P}_{\ell m}(k) \equiv \hat{a}_{lm}(k) \hat{a}_{lm}(k) - \int d^3z \frac{\psi^2(\vec{z})}{\bar{n}(\vec{z})}, \quad (6)$$

where we subtract out the expected shot noise contribution. The light cone projection of the correlation function for flat space (hyperbolic, or “open” models will be covered in a more detailed paper de Laix & Starkman (1998)) has been calculated by Nakamura, Matsubara, & Suto (1997). In terms of a spherical harmonic basis, ξ may be expressed as

$$\begin{aligned} \xi(\vec{z}, \vec{z}') &= \sum_{\ell m} \int \frac{2k^2 dk}{\pi} \bar{D}(z) \bar{D}(z') Y_{\ell m}(\Omega_z) Y_{\ell m}(\Omega_{z'}) \\ &\times \left\{ j_\ell(kr) - \frac{\beta(z)}{k^2} \left[\frac{H(z)}{1+z} \left(\frac{k\ell}{2\ell+1} j_{\ell-1}(kr) - \frac{k(\ell+1)}{2\ell+1} j_{\ell+1}(kr) \right) \right] \right\} \end{aligned} \quad (7)$$

$$- \left. k^2 (A_\ell j_{\ell-2}(kr) - B_\ell j_\ell(kr) + C_\ell j_{\ell+2}(kr)) \right] \Bigg\} \Bigg\{ z \rightarrow z' \Bigg\},$$

with

$$\begin{aligned} A_\ell &= \frac{\ell(\ell-1)}{(2\ell+1)(2\ell-1)} \\ B_\ell &= \frac{\ell^2}{(2\ell+1)(2\ell-1)} - \frac{(\ell+1)^2}{(2\ell+1)(2\ell+3)} \\ C_\ell &= \frac{(\ell+1)(\ell+2)}{(2\ell+1)(2\ell+3)}. \end{aligned}$$

The proper radial distance r is

$$r = \int_0^z \frac{dz}{H(z)}, \quad (8)$$

where we introduce the redshift-dependent Hubble parameter for flat space

$$H(z) = H_0 \sqrt{\Omega_0(1+z)^3 + \lambda_0}. \quad (9)$$

The parameters Ω_0 and λ_0 represent respectively the present day matter density and present day cosmological constant, in units of the critical density, and $\Omega_0 + \lambda_0 = 1$ for flat space. We also use the linear growth factor

$$D(z) = \frac{5\Omega_0 H_0^2}{2} H(z) \int_z^\infty \frac{1+z'}{H(z')^3} dz', \quad (10)$$

along with the linearly evolving bias parameter which, following Fry (1996), can be written

$$b(z) = 1 + \frac{D(0)}{D(z)}(b_0 - 1), \quad (11)$$

to define an overall linear growth factor for the galaxy perturbations relative to the underlying density perturbations:

$$\bar{D}(z) = \frac{D(z)b(z)}{D(0)b(0)}. \quad (12)$$

Finally, we use linear velocity distortion parameter β given by

$$\beta(z) = -\frac{d \ln D(z)}{d \ln(1+z)} \frac{1}{b(z)}. \quad (13)$$

The reader is reminded that eq. (7) is only valid when $\Omega_0 + \lambda_0 = 1$, otherwise an expression with the generalized spherical Bessel functions derived from a hyperbolic Laplacian are required.

Using our expression for the correlation function, we can show that the ensemble average of the estimator in eq. (6) is equal to

$$\langle \hat{P}_{\ell m}(k) \rangle = \int k'^2 dk' P(k') \frac{2}{\pi} \sum_{\ell' m'} I_{\ell' m'}^2(k'), \quad (14)$$

with

$$\begin{aligned}
 I_{\ell m}(k) &\equiv \int d^3z \psi(\vec{z}) \bar{D}(z) Y_{\ell m}(\Omega_z) \\
 &\times \left\{ j_{\ell}(kr) - \frac{\beta(z)}{k^2} \left[\frac{H(z)}{1+z} \left(\frac{k\ell}{2\ell+1} j_{\ell-1}(kr) - \frac{k(\ell+1)}{2\ell+1} j_{\ell+1}(kr) \right) \right. \right. \\
 &\quad \left. \left. - k^2 (A_{\ell} j_{\ell-2}(kr) - B_{\ell} j_{\ell}(kr) + C_{\ell} j_{\ell+2}(kr)) \right] \right\}
 \end{aligned} \tag{15}$$

In the special case of an all sky survey, where $\psi(z) \propto Y_{\ell_o m_o}(\Omega_z)$, one can perform the angular integral $d\Omega_z$ trivially. The estimator for the power in a particular mode is then reduced to the convolution of the true continuum power spectrum with a window function

$$W_{\ell k} = k^2 I_{\ell}^2(k), \tag{16}$$

which is independent of m . It is these windows which indicate how well a power spectrum derived from a limited redshift survey will correspond to the underlying continuum power spectrum.

To demonstrate how light-cone effects will influence the observational window functions defined above, we compare the windows which arise when all redshift effects are included to those obtained ignoring light-cone effects by taking the distance r proportional to the redshift ($r = H_0^{-1}z$) and holding the other redshift dependent parameters — β , H , and $\bar{D}$ — fixed at their $z = 0$ values. Often one sees results calculated in the latter naive limit. For simplicity we consider only volume-limited all-sky surveys, investigating two survey depths: $z_{max} = 0.25$ and $z_{max} = 0.5$, which correspond to the approximate depths of the SDSS magnitude-limited galaxy survey and the SDSS bright red galaxy survey. We take unbiased ($b_0 = 1$), $\Omega_0 = 1$ cosmologies, and consider two values of k , 0.033 and $.33 \, h \, \text{Mpc}^{-1}$.

In figure 1, we show a plot of the naive window functions for $k = 0.033 \, h \, \text{Mpc}^{-1}$ with a range of ℓ 's in a $z < 0.25$ volume-limited survey, ignoring the light-cone effects, *i.e.* this shows the optimal results one can hope to achieve for a volume-limited survey of the specified depth. In figure 2, we show the same results for a realistic survey, including light-cone effects. In figures 3 and 4, we consider the same survey geometry, but calculate the window functions for $k = 0.33 \, h \, \text{Mpc}^{-1}$. The former plot shown the naive window function while the latter shows the realistic result. Next we examine the deeper survey geometry which extends to a depth of $z = 0.5$. Again we consider first the larger scale, $k = 0.033 \, h \, \text{Mpc}^{-1}$, in the naive and realistic limit in figures 5 and 6 respectively. Finally, we examine the smaller scale in figures 7 and 8.

The results shown in the figures described above reveal the importance of considering light-cone effects when estimating power from galaxy surveys, with several effects meriting closer attention. The most striking

is the broadening of the realistic window functions compared to their idealized counterparts. This is caused predominantly by the deviation from the linear distance–redshift relation. At higher redshifts, the larger Hubble expansion rate causes the galaxies to appear more spread out than they do locally. Thus a wave that probes a given scale locally, probes a smaller scale (larger k) at high redshift. A mode in conformal space will then couple to a range of scales when mapped into redshift space, which leads to a broadening in the window. It also causes a readily observed shift in the peak to larger scales. We can estimate the broadening quite easily if we consider the effective frequency at which a conformal mode oscillates in redshift space. For large values of the argument, a spherical Bessel function can be approximated $j_\ell(kr) \approx \sin(kr + \ell\pi/2)/kr$. In a small region Δz located at z , we can rewrite this as $\sin(kr(z) + k\partial r/\partial z \Delta z + \ell\pi/2)/kr(z)$, which has an effective frequency of oscillation of $k\partial r/\partial z$. Recall from eq. (8) that $\partial r/\partial z = H^{-1}(z)$. This tells us that the effective frequency of the conformal modes which describe the density fluctuations will drift from an initial value of k to $H^{-1}(z_{max})k$ in redshift space, so the inverse of this quantity should be a good estimate of the actual window width. For our examples of $z_{max} = 0.25$ and 0.5 , we see that the window widths should be about $1.4k$ and $1.8k$ respectively, which are in good agreement with the results presented in the figures. A final effect, which is less visible, is an overall decrease in the amplitude of the window function. Since the power spectrum decreases with growing z , the overall power measured in a given mode will be less than its local value.

These results are important for analyzing observations, because they indicate that it is impossible to make a clean measurement of the power spectrum, independent of cosmology, from a deep redshift survey. The light–cone effects result in significant broadening and shifting of the window functions when compared to their idealized limit, in a way that depends sensitively on the cosmology. One must deconvolve these effects to compare observations to theory, but this requires knowledge of the underlying cosmological model (perhaps soon to be provided by cosmic microwave background observations). An alternative approach might be to do a Bayesian analysis assuming different cosmologies as priors. One maps the observed data into conformal coordinate space, utilizing the selected cosmology, calculates a power spectrum estimate, and compares the estimate to theoretical predictions based on simple parameterizations of the underlying power spectrum from theories like inflation or defects. For the correct cosmological model, the windows will have minimum width, while the incorrect models will have broadened windows. It is possible that these effects can be used to learn about the underlying cosmology, and even to break the degeneracy in the power spectrum for models in which the shape parameter, Ωh , is constant (we are currently investigating this problem).

Acknowledgments

G.D.S. was supported by an NSF career grant.

REFERENCES

- de Laix, A. A. and Starkman, G. D., in preparation
- Fry, J. 1996, ApJ, 461, L5
- Gunn, J. E., & Weinberg, D. H., in “Wide Field Spectroscopy and the Distant Universe”, ed. S. J. Maddox & A. Aragón-Salamanca (Singapore: World Scientific); Strauss, M. A. 1997, preprint astro-ph/9610032
- Heavens, A. F. and Taylor, A. N. 1995, MNRAS, 278, 73
- Nakamura, T. T., Matsubara, T., and Suto, Y. 1997, preprint, astro-ph/9706034
- Peebles, P. J. E. 1980, The Large Scale Structure on the Universe (Princeton: Princeton University Press)
- Taylor, K., Gray, P. M. 1994, Proc. SPIE 2198, 136, “Instrumentation in Astronomy VIII”, ed. Crawford, D. L. & Craine, E. R.
- Tegmark, M. 1995, ApJ, 455, 429

Figure Captions

Fig. 1.— The naive window function (no light-cone effects) for an all-sky survey of depth $z = .25$ used to optimally estimate the power in the $.033 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 2.— The true, light-cone corrected window functions for an all-sky survey of depth $z = .25$ used to optimally estimate the power in the $.033 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 3.— The naive window function (no light-cone effects) for an all-sky survey of depth $z = .25$ used to optimally estimate the power in the $.33 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 4.— The true, light-cone corrected window functions for an all-sky survey of depth $z = .25$ used to optimally estimate the power in the $.33 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 5.— The naive window function (no light-cone effects) for an all-sky survey of depth $z = .5$ used to optimally estimate the power in the $.033 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 6.— The true, light-cone corrected window functions for an all-sky survey of depth $z = .5$ used to optimally estimate the power in the $.033 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 7.— The naive window function (no light-cone effects) for an all-sky survey of depth $z = .5$ used to optimally estimate the power in the $.33 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.

Fig. 8.— The true, light-cone corrected window functions for an all-sky survey of depth $z = .5$ used to optimally estimate the power in the $.33 h \text{ Mpc}^{-1}$ mode. The results for number of different ℓ modes are shown.















